

TABLE OF CONTENTS

CHAPTER 1 INTRODUCTION AND CELL	1
CHAPTER 2 EPITHELIUM	15
CHAPTER 3 CONNECTIVE TISSUE	29
CHAPTER 4 MUSCLE TISSUE	43
CHAPTER 5 CARTILAGE AND BONE	61
CHAPTER 6 NERVE TISSUE	85
CHAPTER 7 PERIPHERAL BLOOD	107
CHAPTER 8 HEMATOPOESIS	113
CHAPTER 9 CARDIOVASCULAR SYSTEM	127
CHAPTER 10 LYMPHOID SYSTEM	157
CHAPTER 11 SKIN	181
CHAPTER 12 EXOCRINE GLANDS	193
CHAPTER 13 ENDOCRINE GLANDS	205
CHAPTER 14 GASTROINTESTINAL TRACT	223
CHAPTER 15 LIVER AND GALL BLADDER	247
CHAPTER 16 URINARY SYSTEM	261
CHAPTER 17 RESPIRATORY SYSTEM	289
CHAPTER 18 FEMALE REPRODUCTIVE SYSTEM	305
CHAPTER 19 MALE REPRODUCTIVE SYSTEM	329
CHAPTER 20 ORGANS OF SPECIAL SENSE	343
Index	359

This atlas is a series of photographs ranging from low to high magnifications of the individual tissue specimens. The low magnification images should be used for orientation, while the higher magnification images show details of cells, tissues, and organs. Although every effort has been made to faithfully reproduce the colors of the tissues, a full appreciation of histological structure is best achieved by examining the original specimens with a microscope. This atlas is a preview of what should be observed.

The photomicrographs found in this atlas come from the collection of microscope slide used by medical, dental and undergraduate students of histology at the University of Minnesota. Most of these slides were prepared by Anna-Mary Carpenter M.D., Ph.D. during her tenure as Professor in the Department of Anatomy (University of Minnesota Medical School).

Each tissue specimen, in its entirety, has been digitized with a high resolution 40X or 60X lens to generate virtual microscope slides. The Virtual Microscope Collection includes additional slides which complement and extend the core slide collection. Producing the virtual slide collection and developing the web site for their presentation was done with the very capable assistance of Todd C. Brelje Ph.D.

The drawings that appear in the atlas are the product of Jean E. Magney, who is accomplished both as an histologist and an artist. Her talented interpretation of biological structure and its artistic rendering greatly facilitate the learning and comprehension of histology. These drawings first appeared in "Color Atlas of Histology" Stanley L. Erlandsen and Jean E. Magney, Mosby 1992.

Robert L. Sorenson, Ph.D.



First edition 2004 Second edition 2008 (second printing 2011)

INTRODUCTION:

What is histology? Histology is the study of cells, tissues and organs as seen through the microscope. Although this atlas is a guide to biological structure that can be observed through the light microscope, histology also includes cellular detail down to the molecular level that can be observed using an electron microscope. The importance of histology is that it is the structural basis for cell, tissue and organ biology and function (physiology) and disease (pathology).

What is the plan for the study of cells, tissues and organs? Histology is organized into four basic types of tissues.

- 1. Epithelium
- 2. Connective tissue including

Cartilage and bone

Blood and blood formation

- 3. Muscle
- 4. Nervous tissue

Chapters 2-8 are concerned with the features of the four basic tissues. The remaining chapters focus on features of organs. Organs are typically made up of more than one type of tissue and cells with varying degrees of differentiation.

The light microscope, tissue preparation, limits and challenges.

The bright field light microscope is a two lens compound optical instrument. The two lenses are the objective and the oculars. The oculars have a 10 fold magnification and the objectives range from 10X, 20X, 40X to 100X. Thus the total magnification typically ranges from 100 fold to 1000 fold. In practice this means that while using the 10X objective you have a wide field of view, but with low resolution. While using the 100X objective you have high resolution, but with a very small field of view. To use a metaphor what this means is that when using the low power objective you can see the forest but not the trees and while using the high power objective you can see the leaves on the trees but not the forest. Therefore when examining a specimen it is essential to start with the low power objective to gain perspective and then work up to the highest power magnification as needed to observe the necessary detail.

Examination of tissues requires that they be prepared for viewing with a microscope. This is a multi-step process that includes fixation (preserves the tissue), embedment (stabilizes the tissue for sectioning), sectioning (cuts the specimen into thin slices of about 5 um) then placing the sections on a glass slide so they can be stained for viewing.

A note about resolution and detection. Resolution refers to the ability to discriminate between two adjacent objects. For the light microscope with optimal lenses and sample preparation this approaches 0.2 um, which is the theoretical limit for light microscopes. [The eye can resolve about 250-500 um and the electron microscope can resolve about 1 nm) Detection refers to the ability to detect something and this can be much smaller than the limit of resolution. For fluorescence molecules this can be as little as a few molecules!

Structure	Size	Light Micro- scope
Human ovum	120 um	
Most cells	10-30 um	
Red blood cell (RBC)	7 um	
Mitochondium	0.4-1.0 um	
Cilium	0.3 um	
Microvillus	100 nm	Electron Microscope
Microvillus Microtubule	100 nm 24 nm	
	24 nm	
Microtubule	24 nm 15 nm	
Microtubule Myosin filaments Intermediate fila-	24 nm 15 nm	
Microtubule Myosin filaments Intermediate filaments Plasma mem-	24 nm 15 nm 10 nm	

There are several challenges in learning histology. The first being that the view observed through a microscope gives you a perspective that you are unlikely to have experienced previously. It is a complex data set — one with a broad range of shapes and sizes, with varying shades of red and blue. This complex image offers very few clues that are intuitive. Also, the tissue specimen is a two dimensional slice of a complex three dimensional

structure. So, once the two dimensional image has been ascertained you still have the challenge of imagining its three dimensional elaboration. The ideal situation is to have the student and teacher viewing the same specimen simultaneously such as in a dual view microscope. Since this is not always possible, this atlas was written as if a teacher was always at your side to help guide you from low power to the highest power necessary to observe the essential features of the tissue specimens. Thus you will notice that images of all of the slides range from a macroscopic view of the microscope slide itself and then progress through higher magnifications as needed.

How to study microscope slides:

- 1. Know what structures are important to learn. This atlas shows and identifies the structures and how to find them.
- The next task in learning is to see if you can identify the structures when examining a slide. Always start at the lowest power (this is important for context and orientation). Increase the magnification as needed so that additional features of the specimen can be observed.
- 3. Take notes on the features that are observed in the slide. This is best done by drawing pictures and writing a description of the specimen. As in any science laboratory, it is essential that observations be recorded. Not only is this good practice but in research and medicine it is also a legal requirement.
- Each chapter has a section "Observe and note". This lists the features that are essential to learning histology and are noteworthy.

How to take Histology Laboratory Notes:

- A. Draw a picture of the object of interest. (A blue and red pencil is sufficient for nearly all drawings)
- B. Write notes about its appearance, characteristics and features.

Nearly every cell can be described by taking note of:

- 1. Size
- 2. Shape
- 3. Nuclear size and shape and nuclear/cytoplasmic ratio
- 4. General Staining properties (H&E)
 - a. Basophilia & eosinophilia
 - b. Hetero- and euchromatin
- 5. Special staining properties
 - a. Verhoeff, Azan, silver etc.
- 6. Cellular specializations
 - a. Microvilli, cilia, secretion granules, myofilaments etc.
 - b. Unusual amounts of mitochondria, RNA etc
- Cellular constituents such as secretion granule contents (hormones, enzymes)
- 8. Polarity
- 9. Extracellular material
 - a. Extent
 - b. Appearance

10. Location

- a. Example
 - i. Adjacent to similar cells
 - ii. Borders a lumen
 - iii. Surrounded by extensive extracellular matrix
 - iv. Etc.
- 11. Organization (cells, tissues and organs)
 - a. Arrangement of cells of similar and different types
 - b. Arrangement of cells with respect to extracellular material
- 12. Compare and contrast with similar/different cells.
- 13. Heterogeneity among homologous cells:
 - a. Cell development and differentiation
 - b. Cell Cycle
 - c. Active and resting cycles
 - d. Exposure to a concentration gradient of nutrients
 - i. Example
 - 1. Skin cells
 - 2. Liver hepatocytes
- C. Include questions in the notes.

Carefully formulated questions can often reveal the answer.

D. Drawing (and taking notes) is a way of thinking, seeing and understanding.



vi

1

CHAPTER 1 INTRODUCTION AND CELL

The first chapter is an exercise in examining histological tissue specimens through the microscope. A variety of cells, tissues and organs are provided as samples. Also, several different histological stains are used.

A note about stains:

Biological material is inherently of low contrast and provides little to see in the standard brightfield microscope unless treated with a histological stain. 5. Verhoeff: stains elastic protein black

6. Feulgen: stains DNA

7. Sudan: stains lipids

8. PAS (periodic acid Schiff): stains glycogen

9. Aldehyde fuchsin: stains insulin, mast cell granules and elastic fibers purple

HEMATOXYLIN AND EOSIN (H&E):

- 1. This is the most commonly used stain for histology and histo-pathology.
 - a. Hematoxylin: Cationic, positively charged, blue dye complex.
 - i. Reacts with negatively charged groups:
 - 1. COO-- in proteins
 - SO₄ in proteoglycans (GAGs)
 - 3. PO₄ in nucleic acids
 - ii. Reacts with basophilic structures: basophilia
 - Eosin: Anionic, negatively charged red dye
 - i. Reacts with positively charged groups:
 - 1. NH₃+ in proteins
 - 2. mitochondria
 - ii. Reacts with acidophilic structures: acidophilia, eosinophilia

OTHER STAINS:


- 1. Azan: stains collagen blue
- Silver: stains reticular fibers (collagen type III) black
- 3. Golgi stain: stains Golgi apparatus
- Toluidine blue: stains RNA blue and stains mast cells purple

OBSERVE AND NOTE:

- 1. Staining characteristics of various cells and tissues.
- Cell size using red blood cells (7um) as an internal metric.
- Cell size, shape, nuclear size and shape and nuclear/cytoplasmic ratios.
- 4. Eosinophilia and basophilia.
- 5. Euchromatin and heterochromatin.
- 6. Cellular versus extracellular material such as collagen.
- 7. Characteristics of different histological stains.


Cell structure & staining

EM LM





Composite Cell


Nucleus Structure (Euchromatin & Heterochromatin)




Slide # 1 Nuclear morphology & cell size (H&E)








Slide # 2 Cells and Tissues: Size, Shape and Color (H&E)




Slide # 2 Cells and Tissues: Size, Shape and Color (H&E)




Slide # 2 Cells and Tissues: Size, Shape and Color (H&E)





Slide # 2 Cells and Tissues: Size, Shape and Color (H&E)



Slide # 3 (Toluidine Blue)










Slide # 9 Pancreas (Golgi Stain)




Slide # 9 Epididymis (Golgi Stain)







Slide # 12 Feulgen Stain for DNA



CHAPTER 14 GASTROINTESTINAL TRACT

The gastrointestinal tract is a hollow muscular tube that starts at the esophagus and ends with the anus. It is divided into four regions, the esophagus, stomach, small intestine and large intestine. The esophagus is a passage for transporting food to the stomach. The stomach adds gastric juices to begin digestion. It is divided into three histologic regions: cardiac, fundus/body and pyloric. The small intestine is the principle site for digestion and absorption. It transfers chyme from the stomach to the large intestine and is divided into three regions: duodenum, jejunum and ileum. The large intestine has the main function of re-absorbing water from the chyme and adding mucus to facilitate transport of the feces. The parts of the large intestine are the cecum, appendix, colon, rectum and anal canal.

GENERAL PLAN FOR HOLLOW TUBULAR ORGANS

The walls of hollow organs have four layers or tunics: mucosa, submucosa, muscularis externa and adventitia or serosa.

Mucosa (mucous membrane): Mucous membranes line internal passages and provide a barrier between the tissues of the body and the external environment. The membranes are constantly wet and lubricated by mucus. The mucosa has three parts: an epithelium, lamina propria and muscularis mucosa. The epithelium varies in different regions depending on its function (i.e. protective, secretory or absorptive). The lamina propria is a connective tissue layer that supports the epithelium and contains small arteries, veins, lymphatics and nerves. Lymphocytes and plasma cells are also frequently seen in this layer. When glands are found in this layer they are referred to as mucosal glands. The muscularis mucosa, when present, consists of two or three layers of smooth muscle. It facilitates localized movement of the mucous membrane, aiding expression of secretions and movement of fluid across the surface of the epithelium.

Submucosa: The **submucosa** is a layer of fibroelastic connective tissue that supports the mucosa. Found in this layer are blood and lymphatic vessels and nerves. Parasympathetic ganglia found in this layer are called **Meissner's submucosal plexus**. When glands are found in this region (esophagus and duodenum) they are referred to as sub-mucosal glands.

Muscularis externa: This is a separate layer not to be confused with muscularis mucosa. The **muscularis externa** consist of two thick layers of smooth muscle — and inner circular layer and an outer longitudinal layer. Between the layers is a

vascular plexus and an autonomic nerve plexus associated with small parasympathetic ganglia of (Auerbach's) myenteric plexus. The muscularis externa maintains tonus in the tube and propels luminal contents by peristalsis.

Adventitia or serosa: This outermost layer is dense irregular connective tissue. When it blends with connective tissue of the surrounding area it is an adventitia. If it has a free surface projecting into the peritoneal cavity it is covered with a single layer of mesothelial cells (epithelial cells derived from mesoderm) and is called a serosa.

ESOPHAGUS

The epithelium is stratified squamous and non-keratinized. This is a thick layer of 40-60 cells measuring 300-500 um. This is supported by a lamina propria. A well developed muscularis mucosa is present (200-300 um) and surrounded by the submucosal region. Submucosal mucous glands are scattered in this region. The muscularis externa consists of an inner circular layer and an outer longitudinal layer. In the upper third of the esophagus the muscularis is skeletal muscle. In the middle third both smooth and skeletal muscle is present and in the lower third only smooth muscle is present. The myenteric plexus of nerves and ganglia (Auerbach's plexus) are found between the inner and outer layers of the muscularis externa. A tunica adventia is present.

Sтомасн

An abrupt transition occurs at the cardio-esophageal junction, where stratified squamous epithelium gives way to simple columnar epithelium. The simple columnar epithelium (surface mucous cells) dips into the lamina propria to form gastric pits (150-300 um deep). Gastric glands (simple tubular branched) empty into the bottom of the gastric pits. The base of gastric glands rests on a muscularis mucosa. The submucosa is quite prominent and contains numerous arteries, veins, lymphatics and nerves. In the stomach the muscularis externa consists of three layers: a discontinuous inner oblique layer, then an inner circular layer and an outer longitudinal layer. When the stomach is empty the surface is thrown into folds (rugae).

The stomach is divided into three histological regions (cardiac, body/fundus, pyloric) based on their anatomical location and appearance of the glands. The **cardiac** region of the stomach is a narrow rim of tissue around the esophageal opening. The cardiac glands are short tubular glands that are

coiled at the base. The glands consist mostly of mucus secreting cells. Parietal cells may be found in these glands. The fundus and body make up more than 90% of the stomach and have the same histological appearance. The glands of the body and fundus are straight tubular and have three regions: The upper third is the **isthmus** and empties into the gastric pits, the middle third is the neck and the bottom third is the base. There are five types of cells associated with the glands. Regenerative cells are found at the boundary between the isthmus and the gastric pit. These cells are few in number and not readily distinguished in routine preparations. These cells divide and migrate upwards to replenish the surface mucous cells and downward to replenish the rest of the cells in the gastric glands. Mucous neck cells are found in the isthmus and neck region. These cells are scattered among parietal cells and secrete an acidic form of mucus. Parietal cells are distinctive eosinophillic cells with a centrally located nucleus and secrete hydrochloric acid. The eosinophilia is due to the large quantity of mitochondria in these cells. Some parietal cells are also be found in the base of the gland. The primary cell type in the base is the chief cell which has a basophilic cytoplasm in its basal region. Chief cells secrete pepsinogen and gastric lipase. Gastric enteroendocrine cells are part of the diffuse neuroendocrine system (DNES) are few in number and secrete enteric hormones (these can not be identified with H&E). The pyloric region has short coiled tubular glands that only secrete mucus - chief cells and parietal cells are absent.

SMALL INTESTINE

One of the main functions of the small intestine is nutrient absorption. Specializations for increasing surface area for absorption involve three magnitudes of folds or projections.

- Circular transverse folds (plicae circulares or valves of Kerckring) of the entire mucosa (with a core of submucosa) project permanently into the lumen. The plicae are prominent in the duodenum and jejunum and diminish in the later part of the ileum.
- 2. Villi are projections (evaginations) of the mucous membrane (with a core of lamina propria) into the lumen. The shape of villi varies in the different regions of the small intestine: They start as tall, narrow, finger-like projections in the duodenum and evolve to a short broad leaf-like projection in the distal ileum.
- 3. Microvilli are cytoplasmic projections (1-2

um in length) on the surface of the simple columnar epithelial cells. These surface projections make up the **striate border** of intestinal epithelium.

The second main function of the small intestine is digestion and is dependent on secretions from three types of glands:

- Exocrine glands (liver and pancreas) deliver their secretions (bile and digestive enzymes) into the duodenum by way of the cystic duct and main pancreatic ducts.
- Submucosal glands. Submucosal glands are only found in the duodenum (Brunner's glands). They secrete mucus and resemble the pyloric glands of the stomach.
- Intestinal crypts (glands) are invaginations of the surface epithelium down into the underlying lamina propria.

Cell types found in the intestinal epithelium include:

- Simple columnar epithelium absorptive cells have a microvillus (striate) border and are involved in nutrient digestion and absorption.
- Goblet cells secrete mucin.
- Columnar crypt cells transport secretory IgA
- 4. Paneth cells at the base of intestinal crypts produce antibacterial substances. These cells have very eosinophillic secretion granules due to their content of lysozyme.
- 5. M cells occur in regions where lymphoid nodules abut intestinal epithelium. Here the columnar cells are replaced by the cuboidal to squamous M cells. M cells belong to the mononuclear phagocytic system of macrophages and antigen presenting cells.
- Stem cells are located in the base of the intestinal crypts
- **7. Enteroendocrine cells** (DNES) produce hormones and are not readily distinguished in routine preparations.


The **lamina propria** forms the core of the villi and supports the intestinal glands, is highly vascular and rich in lymphocytes and plasma cells. The **muscularis mucosa** lies at the base of the glands and sends fibers into the core of the villi. The **sub-**

mucosa is irregular fibroelastic tissue with a rich lymphatic and vascular supply. Meissner's submucosal nerve plexus is found in this layer and controls the muscularis mucosa. In the duodenum submucosal glands are found. The muscularis externa, is responsible for peristalsis, and has an inner circular and outer longitudinal layer. Auerbach's plexus of nerves is found between the two muscle layers.

The small intestine is divided into three regions: duodenum, jejunum and ileum. The pyloric stomach transitions to the duodenum at the pyloric sphincter (thick inner circular layer of the muscularis externa). The duodenum is the shortest segment (25 cm) and receives secretions from the liver (bile) and pancreas (digestive enzymes). A distinguishing feature of the duodenum is the presence of submucosal glands (Brunner's glands, their appearance differs from pyloric glands only with respect to where they are located i.e. submucosal vs. mucosal). The jejunum and ileum have a similar appearance. Lymphoid tissue in the lamina propria progressively increases from the jejunum to the ileum. In the ileum, permanent clusters of lymphoid nodules (Peyer's patches) become a prominent feature. Villi become shorter, broader and have increasingly larger lacteals (blind ending lymphoid vessels in the core of villi) in the ileum. Frequency of goblet cells and Paneth cells increases as one progresses from the duodenum to the ileum.

LARGE INTESTINE

The main function of the large intestine is to reabsorb water and to consolidate and transport the fecal mass. The parts of the large intestine are the cecum, appendix, colon, rectum and anal canal. The cecum and colon are histologically indistinguishable. Having no villi, the inner surface is smooth and even. The intestinal glands (crypts of Lieberkuhn) are frequent and closely packed together. The glands are simple straight tubular glands and quite long (>600 um). The two major cell types are simple columnar absorptive cells with striated border and numerous goblet cells. Paneth cells may or may not be present. Enteroendocrine cells may be seen at the base of the crypts. Lymphocytes are common in the lamina propria. The muscularis mucosa is found at the base of the glands. The submucosa is well developed with prominent blood and lymph vessels. Meissner's submucosal nerve plexus is easily seen in the colon. The muscularis externa consists of an inner circular layer and an unusual outer longitudinal layer. The outer layer is gathered into three distinct bundles (taenia coli) that are equally spaced around the gut. Between the muscle layers the numerous ganglia of Auerbach's plexus are seen. The colon is mostly covered by a serosa. The **appendix** is a 4-6 cm blind ending diverticulum descending from the cecum. Its epithelium is similar to the colon, but with fewer goblet cells. The crypts are short (150-250 um) in comparison to the colon. Enteroendocrine cells are found in the base of the crypts. Numerous lymphocytes and nodules are present in the lamina propria. When nodules are present M cells are frequently observed in the epithelium overlying the nodules. The muscularis mucosa is very thin. The muscularis externa is inner circular and outer longitudinal layers of smooth muscle. The appendix is covered by a serosa.



OBSERVE AND NOTE:

TONGUE

- Striated muscle
- 2. Lingual (minor) salivary glands
- 3. Foliate and filiform papillae.
- 4. Taste buds
- Taste pore

ESOPHAGUS

- The mucosa consisting of:
 - a. Epithelium: stratified squamous nonkeratinizing
 - b. Lamina propria (the connective tissue support for epithelium in mucous

membranes).

- c. Muscularis mucosa
- 2. Submucosa consisting of:
 - a. Submucosal glands with ducts passing through the mucosa
- 3. Muscularis externa: inner and outer layers
- Ganglia of Auerbach's (myenteric) nerve plexus, located between the inner and outer muscle layers of the muscularis externa.

CARDIOESOPHAGEAL JUNCTION

- Abrupt transition from stratified squamous non-keratinizing epithelium to a simple columnar epithelium.
- Mucosa

- a. Lamina propria
- b. Muscularis mucosa
- 3. Gastric pits
- 4. Cardiac glands
 - a. Mucous cells
 - b. Parietal cells
- Submucosa
- Muscularis externa
- 7. Adventitia

FUNDIC STOMACH

- Rugae
- 2. Muscularis mucosa
- Muscularis externa
- 4. Gastric pits and glands
- 5. Four (five) types of cells characteristic of the stomach:
 - a. Simple columnar epithelium of the surface
 - b. Mucous neck cells
 - c. Parietal cells
 - d. Chief cells
 - e. Enteroendocrine cells cannot be easily recognized with H&E

PYLORIC STOMACH

- 1. Pits and glands
- 2. Short coiled mucosal glands
 - a. Glands consists primarily of cells that secrete mucus
 - i. Absence of parietal cells

PYLORODUODENAL JUNCTION

- 1. Sphincter
- 2. Pyloric stomach
 - a. Mucosa
 - Gastric pits

- ii. Pyloric mucosal glands
- b. Submucosa
- 3. Duodenum
 - a. Mucosa
 - i. Villi
 - ii. Crypts (glands)
 - b. Submucosa
 - i. Submucosal glands (Brunner's glands)

DUODENUM

- 1. Mucosa
 - a. Villi
 - b. Crypts (glands)
 - c. Surface absorptive cells
 - i. Brush border microvilli
 - d. Goblet cells
 - e. Lamina propria
 - f. Muscularis mucosa
- 2. Submucosa
 - Submucosal glands (Brunner's glands)
- 3. Muscularis externa
 - a. Inner circular and outer longitudinal

JEJUNUM/ILEUM

- Mucosa
 - a. Villi
 - i. Compare villi in duodenum, jejunum and ileum
 - ii. lacteals
 - b. Crypts (glands)
 - c. Surface absorptive cells
 - i. Brush border microvilli
 - ii. Terminal web and terminal bar
 - d. Goblet cells

- e. Paneth cells
- f. Lamina propria
 - i. Lymphocytes and plasma cells
 - ii. Lymph nodules (Peyer's patches)
- g. Muscularis mucosa
- h. Plicae circulares

2. Submucosa

- a. Absence of glands in submucosa
- b. Connective tissue, blood and lymph vessels and nerves
- 3. Muscularis externa
 - a. Inner circular and outer longitudinal
 - b. Auerbach's myenteric nerve plexus

APPENDIX

Mucosa

- a. Absence of villi
- b. Short crypts
- c. Surface absorptive cells
- few goblet cells
- e. Lamina propria
 - Lymphocytes, plasma cells, eosinophils
 - ii. Lymph nodules (in some areas M-cells over lie nodule)
- f. Submucosa
- g. Muscularis externa

COLON

1. Mucosa

- a. Absence of villi
- b. Deep crypts (straight intestinal glands)
- c. Surface absorptive cells (microvillus border, terminal web)
- d. Numerous goblet cells
- e. Lamina propria

- i. Lymphocytes, plasma cells, eosinophils
- f. Muscularis mucosa


2. Submucosa


- a. Numerous blood and lymph vessels
- b. Meissner's submucosal nerve plexus


3. Muscularis externa

- a. Inner circular layer, outer longitudinal layer (taenia coli)
- b. Auerbach's myenteric plexus

Slide # 108 Tongue (H&E)











Taste buds














Organization of Stomach








Slide # 112 Fundic Stomach (H&E) Rugae Muscularis mucosa Muscularis externa Slide # 112 Fundic Stomach (Chief & Parietal cells)




Slide # 113 Stomach Fundic (Parietal and Chief cells)












Intestine Structure and Function




Slide # 118 Duodenum, Jejunum and Illeum (H&E)










Slide # 122 Appendix (H&E)



Slide # 123 Colon (H&E)







INDEX

Α

A-band, 43-44, 48 absorptive cells, 224-225, 227-228 acidophils, 205, 207, 209, 211 acinar pancreas, 195 acini, 194, 196-197, 207 actin, 15, 43, iii adenohypophysis, 205 adipocytes, 29-30, 35, 194, 207 adrenal cortex, 128, 206-207, 216-218 adrenal gland, 128, 206-207, 215-219 adrenal medulla, 206, 216, 219 adventitia, 127-130, 133-136, 142, 144-145, 223, 227, 233, 247, 265, 306, 331 afferent lymphatic, 158-159, 163-164 agranulocytes, 107 aldehyde fuchsin, 1, 198, 207, 221-222 alpha-cells, 206, 221 alveolar cells, 290, 292, 302-303, 307 alveolar duct, 290, 292, 301 alveolar macrophage, 290, 292 alveolar pore, 290 alveolar sac, 290, 292 alveolus, 16, 193, 289-290, 292, 301, 309, 344 ameloblasts, 84 ampulla, 305, 307, 322-323, 329, 331, 338, 346 anterior chamber, 343-345 anterior pituitary gland, 207, 209-211 antigen presenting cells, 157, 181, 224 aorta, 127, 130, 143-145 aortic valve, 127, 151-152 appendix, 223, 225, 228, 243 appositional growth, 61-62 arcuate arteries, 262, 308, 318 arcuate artery, 262, 264, 270, 278, 306, 308, 318 area cribrosa, 261 areolar tissue, 29-30, 35-36 arrector pili, 182-183, 186 arrector pili muscle, 182, 186 arteriole, 127-130, 138, 158-160, 174-177, 179, 247-248, 253, 261-264 artery, 127-129, 133-134, 136-144, 150, 158, 248, 261, 264, 270, 278, 306, 341 articular cartilage, 77, 79-80 atrioventricular node, 127 atrioventricular septum, 127 atrium, 43, 127, 130, 147-148 Auerbach's plexus, 86-87, 96-98, 223, 225, 228, 232, 245 autonomic nervous system, 85-86, 206 axon, 85-86, 90, 100-101, 182 axon hillock, 85-86, 90 azan, 1, 30, 33, 35, azure granules, 107, 113-114

band forms, 114, 119 basement membrane, 6, 15-16, 18, 20, 23, 25, 127-129, 159, 181-182, 193, 205, 261, 263, 289, 295, 299, 307, 329-330, 343 basilar membrane, 346 basophilia, 1, 5, 157, iv basophilic erythroblast, 113-114 basophilic metamyelocytes, 122 basophilic myelocyte, 114 basophilic myelocytes, 122 basophilic normoblasts, 116-117 basophils, 107-108, 111-112, 114, 205, 207, 209, 211 beta-cells, 206-207, 221 bile canaliculi, 247-248, 254-255 bile ducts, 247 bile ductule, 247-248, 253 bladder, 16, 247, 258-259, 261-262, 265, 285-287, blood, 1, 4-5, 15-16, 29, 43, 61-63, 107-109, 112-114, 127-128, B-lymphocytes, 157-159 bone, 29, 61-63, 69-70, 72-82, bony labyrinth, 346 Bowman's capsule, 16, 261, 263-264, 268, 276 Bowman's membrane, 343-344 Bowman's serous glands, 289 Bowman's space, 261, 264 brachiocephalic vein, 142-143 bronchi, 16, 289-292, 303 bronchiole, 289-292, 300 Bruch's membrane, 343-345 brush border, 15, 227, 261

С

В

calcified cartilage, 81 calyx, 281-282 canal of Schlemm, 343-344, 350 canaliculi (bone), 62-63, 74 cancellous bone, 63, 69-70, 77 capillaries, 15, 128, 130, 132, 159, 181, 194, 205-207, 248, 261, 263-264, 307, 330-331, 343 capsule cells, 85-86, 91, 95, 159-160, 276 cardiac muscle, 43-44, 49, 51-53, 55-56, 58-59, 127 cardiac skeleton, 43-44, 55, 61, 127, 130, 151-152, 155 cardioesophageal junction, 226 cardiovascular system, 127, carotid artery, 141-142 cartilage, 5, 29, 43, 61-68, 77, 79-81, 127, 289-291, 296, 298, 303, cartilage histogenesis, 64 cell nest, 61-62, 66 cells of Boettcher, 346 cells of Claudius, 346

cells of Hensen, 346 central arteries, 158 central arteriole, 158-160, 174, 176-177, 179 central canal, 86, 89 central lymphoid organs, 157 central nervous system, 85 central vein, 247-248, 251-254 centroacinar cells, 193-194, 196-197 cerebellum, 9 cervical glands, 306, 308, 324 cervix, 306, 308, 324 chief cell, 224 chief cells, 205, 207, 215, 224, 227, 235, 237 chochlear nerve, 346 chondroblast, 61-62, 64-66, 292 chondrocyte, 5, 61-66 chondrogenic layer, 61-62 chorion, 307 choroid, 343-345, 348, 351 chromaffin cells, 206-207, 219 chromophils, 205-206 chromophobes, 205-207, 209, 211 cilia, 15-16, 20, 25, 295, iv ciliary body, 343-344, 348, 350 ciliary glands, 344-345, 354-355 cilliated cells, 323 clear cells, 181, 205, 207, 215 cochlea, 346, 356 cochlear duct, 346, 356-357 collagen, 1, 5-7, 29-30, 33, 43, 61-62, 86, collecting ducts, 261, 263-265, 282 collecting tubules, 261, 264, 269-271, 273, 279 collecting veins, 128, 178 compact bone, 62-63, 69-70 conjunctiva, 344-345, 353-355 connective tissue, 15, 29-30, 33, 35-38, 43, convoluted tubules, 261, 263-264, 267-268, 274, 278-279 cornea, 343-344, 348 corneal-scleral junction, 343 corona radiata, 305, 313 coronary artery, 150 corpora amylacea, 330-331 corpora cavernosae, 330-331 corpus albicans, 305, 307 corpus cavernosum, 340-341 corpus hemorrhagicum, 305, 307, 315 corpus luteum, 305, 307, 316 corpus spongiosum, 330-331, 340 cortex (adrenal gland), 128, 206-207, 216-218 cortex (kidney), 157, 261-264, 267-268, 270-272, 278 cortex (lymph node), 157-159, 163-164, 166 cortex (ovary), 307 corticotropes, 205 crista ampullaris, 346-347, 358 crypt cells, 224 crypts of Lieberkuhn, 225, 244 cumulus oophorus, 305, 307, 313

360 **INDEX** cuticle, 182 cystic duct, 224 decidua basalis, 307 decidua capsularis, 307 decidua parietalis, 307 dendrite, 85-86, 90 dendritic cells, 157-159 dense irregular connective tissue, 29, 36, 38, 127-128, 223, 247, 305-306, 308, 329, 343 dense regular connective tissue, 29, 36-37, 61, 330 dental papilla, 83 dental pulp, 84 dentin, 84 dermal papillae, 86, 181-182, 188 dermis, 29, 38, 86, 181-182, 184, 189-190, 309 Descement's layer, 343 desmosomes, 15, 157, 181 diastole, 127 diffuse lymphocytic infiltrations, 158 distal convoluted tubule, 261, 263-264, 268, 274, distributing arteries, 128 DNA, 1, 14 dorsal horn, 85-86, 89 dorsal root ganglion, 85-86, 88, 91-93 dorsal roots, 85 duct cells, 6, 183, 193, 197, 307 ducts of Bellini, 261, 264 ductus deferens, 329-331 dust cells, 157, 301-302 Ε ear, 61, 346, 356-358 efferent ductules, 329 efferent lymph vessels, 158 ejaculatory duct, 330 elastic artery, 127-130, 134, 141, 143-144 elastic cartilage, 61-62, 67-68, 289-291, 296 elastic fibers, 1, 29-30, 61, 86, 127-129, 181, 290-291, 306 elastic lamina, 127-130, 133-134, 145 elastic tissue, 29-30, 62, 127-129, 306, 343 elastin, 33-34 endocardium, 127, 130, 148-149 endochondral bone growth, 63 endocrine glands, 15, 128, 205, i endometrium, 306, 308, 317, 320 endomysium, 43-44, 47, 50-51 endoneurium, 85-87, 93, 100

endosteum, 62-63

263-264, 290, 331

eosin, 1, 30

endothelial cell, 5, 15, 127-130, 158-160, 247-248,

enteroendocrine cells, 224-225, 227

eosinophilia, 1, 5, 193, 224, 261, 263, iv eosinophilic metamyelocytes, 122 eosinophilic myelocytes, 122 344 eosinophils, 29-30, 41, 107-108, 111-112, 114, 122, 228 epicardium, 127, 130, 148-149 epidermis, 15, 181-182, 184, 189-190 epididymis, 329, 331, 336 epiglottic cartilage, 289 epiglottis, 61, 67-68, 289, 291, 294-296 epimysium, 43-44 epineurium, 86-87, 99 epiphyseal plate, 63, 77, 79 epithelial reticular cells, 157, 160, 169-171 epithelium, 15-16, 18-20, 22-27, 83, erythropoesis, 113 esophagus, 16, 26, 96-97, 223, 226, 230-232 euchromatin, 1, 6, iv exocrine glands, 15, 193, 224, 290, i exocrine pancreas, 193, 196 external elastic lamina, 128-130, 133-134 external root sheath, 182, 188 extracellular matrix, 29, 343, v extraglomerular mesangium, 263, 272 extralobular ducts, 193-194, 197, 199 eye, 343-344, 348-353, iii Н evelid, 344-345, 354-355 eyelids, 344

F

Fallopian tube, 305, 322-323 false vocal cords, 289 fascicle, 43, 48, 86 fat cells, 4, 7, 29-30, 38, 158, 206, 247 female reproductive system, 305, i fetal liver, 247-248, 258 fetal lung, 292, 304 Feulgen, 1, 14 Feulgen stain, 14 fibria, 322 fibroblast, 5-7, 29-32, 43, 61, 85-87, 129, 182, 247, 263, 290, 305, 329-330, 343 fibrocartilage, 55, 61, 68 fibroelastic lamina propria, 289 fibrogenic layer, 61-62 fimbriae, 305 follicles, 182, 184, 186-188, 205, 207, 213, 305, 307, 311-314, 344-345 follicular cells, 205-207, 214, 305, 307 follicular phase, 305

G

gall bladder, 247, 258-259, i gastric glands, 223-224, 227, 234, 238 gastric pits, 223-224, 227, 234 gastrointestinal tract, 86, 223, i germinal center, 158, 160, 165

germinal epithelium, 305, 329-330 gland cells, 5, 183, 193, 202-204, 207, 214-215, glands of Littre, 331, 341 glans penis, 330 glassy membrane, 182, 188 glia, 85 glomerulus, 261, 263-264 glucagon, 206 glycogen, 1, 43, 127, 247-248, 255-256, 306 goblet cell, 15-16, 20, 24-25, 224-225, 227-228, 241, 244, 289-291, 295, 344 Golgi apparatus, 1, 11, 205 Golgi stain, 1, 10-11 gonadotropes, 205 Graafian follicle, 305, 312-313 granulocytes, 107 granulomere, 107 granulopoesis, 113 granulosa, 305, 307, 312-313, 316 granulosa cells, 305, 307, 316 granulosa luteal cells, 307, 316 grey matter, 85 ground substance, 29, 61-62

hair bulb, 182, 188 hair follicles, 182, 184, 186-188 Hassel's corpuscles, 157-158, 160 Haversian, 62-63, 70-74 Haversian canal, 63, 70-74 Haversian lamellae, 62-63, 70-71, 74 H-band, 43-44, 48 heart, 43, 127-130, 154 heart conduction system, 127 helicine artery, 341 helicotrema, 346, 356 hematopoesis, 113, 247, 258, i hematoxylin, 1, 12-13, 61 Henle's loop, 261, 263, 269, 271, 273, 279, 281 hepatic arteriole, 247-248, 253 hepatic ducts, 247 hepatic sinusoids, 247, 254-255 hepatic stellate cells, 247 hepatocytes, 247-248, 252, v Herring bodies, 205, 207, 210, 212 heterochromatin, 1, 6, 43 high endothelial venules, 158-159, 166 high resistance channels, 128 horny cells, 181 hyaline cartilage, 61-62, 65-66, 289-291, 303 hyalomere, 107 hypodermis, 86, 181-182, 184, 189

ı

I-band, 43-44, 48 ileum, 223-225, 227, 241

immature bone, 63 infundibulum, 305, 307, 322-323 inlet vessels, 247 inner circumferential lamellae, 62, 73 inner enamel epithelium, 83 inner root sheath, 182 insulin, 1, 206-207 interalveolar septum, 290 intercalated disks, 43-44, 52-53 intercalated ducts, 193-194, 200 interlobar vessels, 261, 264 interlobular arteries, 262 interlobular ducts, 193-194, 196-197 internal elastic lamina, 128-129, 133-134 Interstitial cells of Leydig, 329, 331, 333 interstitial growth, 61-62 interstitial lamellae, 62-63, 70, 74 interterritorial matrix, 61-62 intervetebral disk, 68 intestinal crypts, 224-225, 228 intestinal glands, 224-225, 228 intralobular duct, 193, 197 intralobular ducts, 193-194, 196, 200, 309 intramembranous bone formation, 75-76, 82 intramembranous bone growth, 63 iris, 343-344, 348-349 islets of Langerhans, 193-195, 198, 206-207, 220-222 isthmus, 224, 305-306 J

jejunum, 223-225, 227, 239-240 junctional complex, 15, 43, 193 juxtamedulary cortex, 164 juxtamedullary renal corpuscles, 262, 264

Κ

keratin, 15-16, 26-27, 181 keratinocytes, 181 keratinosome, 181 keratohyalin granules, 181 kidney, 15-16, 107, 113, 157, 205, 261-264, 267-282 Kupffer cells, 157, 247-248, 257

L

lacrimal gland, 344, 353 lacteals, 225, 227 lactiferous duct, 306, 326 lactiferous sinus, 307 lactotrope, 205-206 lacunae, 61-63, 70-74, 307 lamellae, 62-63, 70-71, 73-74 lamina propria, 29, 158, 223-228, 231, 247, 262, 265, 289, 291-292, 306, 308-309, 329, 331, 344 Langerhans cells, 157, 181, 207 Large intestine, 223, 225 larynx, 25, 289, 291, 297 lens, 343-345, 348-349, iii Leydig cells, 329, 331, 333 limbus, 343-344, 346, 348 lipofuscin pigment, 43-44, 52, 86, 206, 218 Littre, 330-331, 341 liver, 8, 15, 128, 157, 224-225, 247-248, 250-258, i, v liver hematopoesis, 247, 258 liver lobules, 247-248, 250-251, 253 lobules, 157, 193-194, 247-248, 250-251, 253, 307, 309, 330 loose connective tissue, 29, 127, 129, 158, 307, 329, 331, 343 lung, 15-16, 127, 157, 290-292, 299-304 luteal phase, 305 lymph node, 4, 9, 40-41, 158-159, 162-167 lymph node cortex, 163, 166 lymph node medulla, 163 lymph nodes, 29, 157-158 lymph nodules, 158-159, 228, 289, 291 lymphatic vessel, 139, 158-159, 162, 223, 247-248 lymphocyte nuclei, 4 lymphocytes, 107-108, 110, 112, 124, 157-160, 223-225, 228, 306-307 lymphoid system, 157, i

M

M cells, 224-225 macrophage, 29-32, 41-42, 157-160, 162-163, 224, 247-248, 263, 290, 292, 301, 305-306, 329, macrophages, 157, 159, 224, 247-248 macula, 261, 263-264, 275, 279-280, 344, 346 macula densa, 261, 263-264, 275, 279-280 major calyces, 262 male reproductive system, 329, i mammary gland, 305-306, 309, 325-327 mammotropes, 205 marginal zone, 159-160, 177 mast cells, 1, 29-30, 39-40, 329, 331 mature (Graafian) follicle, 305 mediastinum testis, 329, 335 medium and large veins, 128 medulla, 157-160, 163-164, 182, 206-207, 216, 219, 261-264, 267, 270-271, 278, 305 medulla (adrenal gland), 206-207, 216, 219 medulla (kidney), 157, 261-264, 267, 270-271, 278 medulla (lymph node), 157-159, 163-164, 305 medulla (ovary), 305 medullary cords, 158-159 medullary ray, 261, 263-264, 270, 272-273, 278, 282 medullary region, 157, 261 megakaryocytes, 114, 123

Meibomian glands, 344-345, 354-355 Meissner's corpuscles, 86, 182 Meissner's plexus, 86-87, 97-98, 223 melanin, 181, 189 melanocytes, 181, 183, 188 membranous labyrinth, 346 menstrual phase, 306 Merkle cells, 181 mesangial cells, 157, 263-264, 276 mesangium, 263, 272, 276-277 mesenchymal cells, 30-32, 63 mesentery nerves, 102-103 metamyelocytes, 119-120, 122 microvilli, 15-16, 20, 24, 224, 227, 241, 289, 307, 329, iv minor calvces, 262 mitochondria, 1, 12-13, 63, 193, 205, 224, iv modiolus, 346, 356 monocytes, 63, 107-108, 110, 112, 125, 157 mucoid connective tissue, 29 mucosa, 158, 223-228, 230-231, 233, 235, 238, 243, 247, 262, 289, 305-308, 330 mucous, 5, 15-16, 193-194, 202-204, 223-224, 226-227, 236, 289, 303, 306, 331, 344 mucous cells, 5, 15, 193-194, 202-204, 223-224, 227, 236 mucous glands, 193, 223-224, 227, 303, 306, 331 mucous neck cells, 224, 227, 236 multilaminar primary follicles, 307, 311, 314 muscle, 7, 15, 33, 37, 43-44, 47-53, 55-59, 86, muscle fascicle, 43, 48 muscle insertion, 44, 50-51 muscular artery, 129, 133-134, 136-140 muscularis externa, 223, 225-228, 230-233, 235, muscularis mucosa, 223-228, 230-231, 233, 235, 238, 243, 247, 305-306 myelin, 85-87, 100-101 myeloblast, 113-114, 119-120 myelocytes, 119-120, 122 myenteric plexus, 223, 226, 228, 232 myocardium, 127, 130, 149 myoepithelial cells, 193, 307, 343-344 myofibrils, 43-44, 50, 127 myometrium, 306-308, 317, 320 myoneural junctions, 43 myosin, 43, iii myo-tendinous insertion, 43-44

Ν

nasal cavity, 289, 291 nasopharynx, 289 nephron, 261, 263 nerve, 43, 85-87, 99-101, 128, 182, nerve fascicles, 87, 99 neurohypophysis, 205 neuron, 85-86, 289 neutrophil, 113-114 neutrophilic band, 113-114, 120-121 neutrophilic band cells, 121 neutrophilic metamyelocyte, 113-114, 120 neutrophilic myelocyte, 113-114, 120 neutrophilis, 29, 107-108, 112-114, 159 Nissl bodies, 86, 95 Nissl substance, 4, 85-86, 90 node of Ranvier, 86-87, 100-101 nodule, 158-159, 164, 172, 174, 177, 179, 228 non-striated muscle, 43 nucleolus, 4, 9-10, 85, 329 nucleus, 4, 9-10, 29, 43-44, 85-86,

0

odontoblasts, 84 olfactory region, 289 oocytes, 305, 307 optic disk, 344-345, 352 optic nerve, 343-345, 352 ora serrata, 343, 345 oral epithelium, 83 organ of Corti, 346, 357 orthochromatic erythroblast, 113-114 orthochromatic normoblasts, 116-117 osseous spiral lamina, 346 ossified cartilage, 80 osteoblasts, 61-63, 76, 78, 81-82 osteoclasts, 63, 78-79, 81 osteocyte, 62-63, 70-74, 76, 82 osteocyte lacunae, 63, 70-74 osteoid, 62-63, 76, 82 osteon, 62-63 otoliths, 346 outer circumferential lamellae, 62-63, 73 outer enamel epithelium, 83 ovarian cycle, 305-306 ovary, 16, 305, 307, 311-316 oviducts, 305-306 ovulation, 305 oxyphils, 205-207, 215

Ρ

Pacinian corpuscle, 86-87, 104-105, 182, 192 palate, 291, 293-294 palatine tonsil, 172 palpebra, 344, 353-354 pancreas, 10, 16, 39, 128, 193-198, 206, 220-222, 224-225 pancreatic ducts, 16, 224 Paneth cells, 224-225, 227-228, 242 papillary layer, 181 paracortex, 158-159 parafollicular cell, 205, 207, 214 parasympathetic ganglion, 86-87, 96-98, 232 parasympathetic nerves, 86, 223 parathyroid gland, 205-207, 214-215 paratrabecular sinuses, 158

pupil, 343-344, 348

parietal cells, 224, 227, 235, 237 parotid gland, 193-194, 199-201 pars basalis, 308, 317, 321 pars distalis, 205 pars functionalis, 306, 308, 317 pars intermedia, 205 pars tuberalis, 205 PAS, 1, 43, 54-55, 127, 193, 247-248, 255-256 pasasympathetic ganglion, 97 pectinate muscle, 147 penicillar arteries, 159 penis, 329-331, 340-341 periarteriolar lymphocytic sheath, 158, 174, 179 peribiliary capillary, 253 perichondrium, 61-62, 64-66 pericyte, 128 perimysium, 43-44 perineurium, 86-87, 99 periodic acid Schiff, 1, 247 periosteum, 37, 51, 62-63, 79, 82 peripheral lymphoid tissues, 158 peripheral nerve, 85, 87, 99-101 peripheral nervous system, 85 Peyer's patches, 158, 225, 228, 242 pituicytes, 205, 207 pituitary gland, 205, 207-212 placenta, 305, 307, 309, 328 plasma cells, 29-30, 40, 114, 124, 157-159, 223-224, 228, 307 platelets, 107-108, 114 plicae circulares, 224, 228 pneumocytes, 290, 292, 302 podocytes, 261, 263-264, 276-277 polychormatic normoblasts, 116 polychromatic erythroblast, 113-114 portal canal, 247-248, 252-253 portal venule, 247-248, 253 post capillary venules, 128-129 posterior chamber, 343, 345 posterior pituitary gland, 205, 210, 212 PP-cells, 206 predentin, 84 primary nodule, 158-159, 164 primary nodules, 158-159 primary oocytes, 305, 307 primordial follicles, 305, 307, 311 proerythroblast, 113-114 proliferative phase, 306 promyelocyte, 113, 119-120 promyelocytes, 119-120 pronormoblasts, 116 prostate, 330-331, 339-340 proximal convoluted tubule, 261, 263-264, 267-268, 274, 278-279 pseudostratified columnar epithelium, 15-16, 20, 25, 289, 295, 297, 299, 303, 329-331, 344 pseudounipolar, 85 PTA stain, 49, 53, 56 pulp arteriole, 159, 174-177

Purkinje fibers, 44, 54, 56, 127, 130, 152-154 R radial arteries, 306, 308, 318 Rathke's cysts, 205, 207-208 RBC, 4, 7, 113, iii reaction center, 174 red blood cells, 1, 5, 107-108, 158-159, 264 red pulp, 158-160, 173, 175-177 regenerative cells, 224 Reissner's vestibular membrane, 346 renal arteries, 262 renal columns, 261-262 renal corpuscle, 261-264, 267-268, 274-277, 279 respiratory bronchioles, 289-291, 300 respiratory epithelium, 16, 289, 291, 299 respiratory system, 15, 289, i resting zone, 63, 78, 80 rete testis, 329, 331, 335 reticular fibers, 1, 29-30, 34, 43, 86, 128-130, 132, 158-159, 167, 181, 247, 290, 329 reticular layer, 181-182 reticular tissue, 29-30, 329 reticulocyte, 113, 116 retina, 343-345, 348, 351-352 retina layers, 345, 352 RNA, 1, 4, 8-10, 85, iv root sheath, 182, 188 rugae, 223, 227, 235, 247 S saccule, 346-347 satellite cells, 85 scala media, 346 scala tympani, 346, 356-357 scala vestibuli, 346, 356-357 scalp, 182, 184-188 Schwann cell, 85-87, 100-101 sclera, 343-344, 348, 351 sebaceous gland, 182-183, 185-186, 344-345 secondary follicles, 305, 307, 312-313 secondary nodule, 158-159, 164 secretion granules, 193, 205, 224, iv secretory phase, 306 semicircular canals, 346-347, 356 seminal vesicle, 329-331, 338-339 seminiferous germinal epithelium, 329 seminiferous tubules, 329-330, 332, 334 sero-mucous glands, 289, 291 serosa, 223, 225, 247, 305-306 serous cells, 193-194, 202-204, 289-290, 344 serous demilune, 203 serous glands, 193, 289 serous secretory cells, 193

Sertoli cells, 329-330, 333-335

sheathed arteriole, 159-160, 175

sheathed capillary, 177 silver stain, 128, 158, 167 simple columnar epithelium, 19, 24, 223-224, 226-227, 241, 247, 289, 306-308, 330, 346 simple cuboidal epithelium, 18-19, 23, 182, 193, 197, 205, 247, 261, 263, 289, 305, 307, 311 simple squamous epithelium, 16, 18, 23, 261, 263, 289, 343 sino-atrial node, 43, 127 sinuses, 158-159, 306, 331 sinusoids, 128, 159-160, 175, 177-179, 206-207, 217, 247-248, 254-255 skeletal muscle, 43-44, 47-51, 53, 58-59, 223, 289, 291, 306 skin, 15-16, 26-27, 29, 86, 103-105, 157, 181-182, 189-192, small collecting veins, 128 small intestine, 223-225 smooth muscle, 7, 43-44, 49, 56-59, 127-129, 182, 223, 225, 247, 262-263, 289-292, 306-307, 309, 329-331, 343-344 soma, 85 somatostatin, 206 somatotropes, 205 space of Disse, 247-248 specific granules, 107, 113-114 spermatids, 329-330, 334 spermatocytes, 329-330, 334 spermatogonia, 329-330, 333-334 spermiogenisis, 334 spinal cord, 4, 85-86, 88-91 spiral ganglion, 346, 356-357 spleen, 128, 157-158, 160, 173-178 splenic artery, 158 splenic cords, 159-160 splenic sinusoids, 159, 175 spongy bone, 62, 69 stellate reticulum, 83 stereocillia, 329, 331, 346 stomach, 16, 193, 223-225, 227, 233-238 straight tubules, 261, 264 stratified cuboidal, 16, 193, 307, 309 stratified squamous epithelium, 25-27, 159, 181-182, 223, 226, 289, 291, 295, 297, 306, 308, 343-344 stratified squamous keratinized epithelium, 181-182, 344 stratum basalis, 181-182, 189-190 stratum corneum, 181, 189-190 stratum germinativum, 181, 190 stratum granulosum, 181, 189-190 stratum intermedium, 84 stratum lucidum, 181 stratum spinosum, 181, 189-190 striate border, 224 striate ducts, 193, 200 striated muscle, 43, 226 subcapsular sinus, 158-159, 165

sublingual gland, 193-194, 203-204

submandibular gland, 193-194, 201-203 submucosa, 223-228, 230, 233, 247 submucosal glands, 223-227, 238, 240, 330 surface mucous cells, 223-224, 227 sweat glands, 182, 184-185, 187, 190, 193, 345 sympathetic ganglion, 86, 88, 91, 95 syncytiotrophoblasts, 307, 328 systole, 127

Τ

taenia coli, 225, 228, 243 tarsal glands, 345, 354-355 tarsal plate, 344-345, 355 tectorial membrane, 346, 357 tendon, 37, 50-51 teritorial matrix, 66 terminal bar, 15-16, 20, 227 terminal bronchioles, 289-291, 300 terminal pulp capillaries, 159 terminal web, 15-16, 20, 227-228, 241 territorial matrix, 61-63 testis, 128, 329-335 theca externa, 305, 307, 313 theca interna, 305, 307, 313 theca luteal cells, 305, 316 thick skin, 27, 86, 103-105, 181-182, 190-192 thin skin, 26, 86, 181-182, 189, 289 thymus, 157, 160, 168-171 thyroid follicles, 207, 213 thyroid gland, 205-207, 213-214 thyrotropes, 205 T-lymphocytes, 157-159 toluidine blue, 1, 8-10 tonofilaments, 181 tonsil crypt, 172 tonsils, 158, 289 tooth development (bell stage), 83-84 trabeculae, 62, 157-160, 330-331 trabecular sinus, 159, 165 trabecular veins, 159-160, 178 trabecular vessels, 158-159, 176, 178 trachea, 25, 65-66, 289-291, 298-299 tracheal cartilage, 289, 298 trachealis muscle, 289, 291, 298 transitional epithelium, 15-16, 22-23, 262, 265, 283-287 tricuspid valve, 150 tubuli recti, 329, 331, 335 tunica adventitia, 127-130, 133-136, 142, 144-145 tunica albuginea, 305, 329-331, 340 tunica intima, 127-130, 133, 135, 144-145 tunica media, 127-130, 133, 135, 142, 144-145 type I pneumocytes, 290, 292, 302 type II pneumocytes, 290, 292, 302

U

umbrella cells, 16, 22-23, 262, 265, 285-287

unilaminar primary follicles, 307, 311, 314 ureter, 16, 261-262, 265, 283-284 urethra, 261, 329-331, 340-341 urinary bladder, 16, 265, 285-287 urinary pelvis, 261 urinary pole, 261, 263-264, 268, 275-277, 280 urinary system, 261, i uriniferous tubule, 261 uterine glands, 306, 308, 317-318 uterine tube, 305, 322-323 uterus, 16, 43, 305-308, 317-321 uterus (late secretory), 308, 319 uterus (menopausal), 321 uterus (menstrual), 308, 320-321 uterus (proliferative), 308, 317-318 uterus (secretory), 308, 319 utricle, 346-347

V

vagina, 16, 305-306, 308, 325 valves of Kerckring, 224 vas deferens, 329, 331, 336-338 vas deferens (ampulla), 331, 338 vasa recta, 262, 264, 270-271, 278 vasa vasorum, 127, 129 vascular pole, 261, 263-264, 275-276, 280 vaso vasorum, 130, 142-143, 145 vein, 128, 135-137, 140-143, 146, 159, 247-248, 251-254 vein valve, 146 vena cava, 130, 145-146 ventral horn, 85-86, 89 ventral motor neurons, 89-90 ventral roots, 85 ventricle, 127, 130, 149, 289, 291 ventricular fold, 289 venule, 130, 138-139, 253 Verhoeff, 1, 30, 35, 67-68, 128, 291, iv vestibular apparatus, 356, 358 vestibular membrane, 346, 356-357 vestibule, 289, 306, 346 villi, 224-225, 227-228, 307, 309 vitreous body, 343 vitreous chamber, 343 vocal cords, 289 vocal fold, 289, 291 vocal ligament, 291, 297 vocalis muscle, 289, 291, 297 Volkman canal, 73 Volkmann's canals, 62

W

Wasserhelle cells, 205, 207, 215 white blood cells, 4-5, 107-108, 128 white pulp, 158, 160, 173-174, 176

Z-line, 43-44, 48 zona fasciculata, 206, 216, 218 zona glomerulosa, 206, 216-217 zona pellucida, 305, 307, 313 zona reticularis, 206, 216, 218-219 zone of calcification, 78, 80-81 zone of hypertrophy, 63, 78, 80 zone of proliferation, 63, 78, 80

Slides

Slide 1 Lymph Node, 4 Slide 1 Nuclear Morphology & Cell Size, 3 Slide 1 Spinal Cord, 4, 90 Slide 3 Toluidine Blue, 8-10 Slide 2 Cells and Tissue: Size, Shape, Color, 5-7 Slide 9 Golgi Stain, 10-11 Slide 10 Iron Hematoxylin Stain, 12-13 Slide 12 Feulgen Stain, 14 Slide 16 Gut: Smooth muscle, 57-58 Slide 16 Simple Epithelia, 18-20 Slide 17 Stratified Epithelia, 21 Slide 18 Transitonal Epithelium, 287 Slide 20 Pig Snout, 31-32, 64, 75-76, 83-84 Slide 20 Pig Snout Embryo, 64, 75-76, 83-84 Slide 21 Connective tissue H&E, 35 Slide 22 Connective tissue Verhoeff, 35 Slide 23 Connective tissue Azan, 35 Slide 23 Tendon: Muscle insertion (Azan), 51 Slide 24 Connective Tissue H&E, 33 Slide 24 Mesentery, 38, 41, 102, 132-139, 162-163 Slide 24 mesentery Lymph Node, 41, 162-163 Slide 24 Mesentery nerves, 102 Slide 24 Vessel: Smooth muscle, 57 Slide 25 Connective Tissue Verhof, 33 Slide 25 Mesentery (Verhof), 33, 134-135, 137-139 Slide 26 Connective Tissue Azan, 33 Slide 26 Mesentery (Azan), 42, 102-103, 134, 136-137 Slide 26 Mesentery nerves (azan), 102-103 Slide 29 Endochondral Bone Formation, 77-79 Slide 29 Muscle Attachment, 37 Slide 29 Tendon: Muscle insertion, 51 Slide 30 Tendon: Muscle insertion, 50 Slide 31 Liver (Trypan Blue), 256-257 Slide 33 Blood Smear, 109, 112 Slide 34 Bone Marrow Smear, 115 Slide 36 Epiglottis (H&E), 67-68 Slide 38 Epiglottis, 294-296 Slide 39 Epiglottis (Verhof), 67-68, 296 Slide 40 Intervetebral Disk: Fibrocartilage, 68 Slide 43 Epiphysis - Cancellous and compact Bone, 69 Slide 43 Skill - Cancellous and compact bone,

69-70

Slide 245 Heart Purkinje Fibers, 154

Slide 44 Ground bone, 73-74 Slide 126 Liver, 250-252 Slide 45 Decalcified bone, 72 Slide 127 Liver (Azan), 253-255 Slide 46 Endochondral Bone Formation, 79-82 Slide 128 Liver (PAS), 255-256 Slide 130 Gall Bladder, 258-259 Slide 46 Intramembranous bone formation, 82 Slide 47 Spinal Cord, 88-89, 91 Slide 131 Pancreas, 195-197, 220 Slide 50 Dorsal Root Ganglion, 91-92 Slide 132 Pancreas (Azan), 198 Slide 51 Dorsal Root Ganglion (Azan), 93 Slide 133 Pancreas (Aldehyde Fuchsin), 198, Slide 52 Peripheral Nerve, 99-101 221-222 Slide 134 Nasal Conchae and Palate, 293 Slide 53 Smooth Muscle, 56 Slide 54 Cardiac Muscle, 51-53 Slide 134 Palate, 293-294 Slide 55 Skeletal Muscle, 47-49 Slide 135 Larynx, 25, 297 Slide 56 Skeletal, Cardiac and Smooth Muscle Slide 136 Trachea, 25, 65-66, 298-299 (PTA stain), 49, 53 Slide 136 Trachea, Hyaline Cartilage, 65-66 Slide 57 Skeletal Muscle Teased, 50 Slide 137 Lung (fetal), 304 Slide 58 Cardiac Skeleton, 55, 155 Slide 139 Lung, 303 Slide 58 Cardiac Skeleton (PAS), 53, 55, 155 Slide 140 Fetal Kidney, 267-269 Slide 58 Purkinje Fibers, 54, 153-154 Slide 141 Kidney, 270-277 Slide 58 Purkinje fibers (PAS), 54 Slide 141 Simple epithelium, 23 Slide 59 Sympathetic Ganglion, 95 Slide 142 Kidney, 278-281 Slide 142 Kidney (Azan), 278-281 Slide 61 Popliteal Artery and Vein, 140-141 Slide 62 Popliteal Artery and Vein, 140-141 Slide 143 Kidney, 281-282 Slide 63 Brachiocephalic Vein, 142-143 Slide 146 Ureter, 283-284 Slide 63 Carotid Artery, 141-142 Slide 147 Bladder, 285 Slide 65 Aorta, 143-145 Slide 149 Pituitary Gland, 208-210 Slide 65 Vena Cava, 146 Slide 150 Pituitary Gland (Azan), 210-212 Slide 66 Aorta (Verhof), 144-145 Slide 151 Thyroid Gland, 213-214 Slide 66 Vena Cava (Verhof), 145-146 Slide 154 Parathyroid Gland, 214-215 Slide 67 Vein Valve, 146 Slide 155 Adrenal Cortex, 218 Slide 69 Pectinate Part of Right Atrium, 147 Slide 155 Adrenal Gland, 215-219 Slide 69 Smooth part of Left Atrium, 148 Slide 155 Adrenal Medulla, 216, 219 Slide 160 Ovary, 311-312 Slide 70 Left Ventricle, 149 Slide 70 Right Ventricle, 149 Slide 165 Uterus (proliferative), 317-318 Slide 71 Coronary Artery, 150 Slide 166 Uterus (late secretory), 319 Slide 71 Tricuspid Valve and Coronary Artery, 150 Slide 167 Uterus (menstrual), 320-321 Slide 72 Coronary Artery (Verhof), 150 Slide 170 Uterus (menopausal), 321 Slide 73 Aortic Valve, 151-152 Slide 171 Fallopian Tube, 322-323 Slide 74 Cardiac Skeleton, 152 Slide 172 Cervix, 324 Slide 74 Purkinje Fibers, 56, 152-153 Slide 173 Vagina, 325 Slide 76 Lymph Node, 164-166 Slide 174 Mammary Gland, 325-327 Slide 78 Reticular fibers, 34 Slide 175 Placenta, 328 Slide 79 Thymus, 168-171 Slide 178 Testis (adult), 333-335 Slide 81 Palatine Tonsil, 172 Slide 178 Testis (neonate), 332-333 Slide 84 Spleen, 176-178 Slide 181 Vas Deferens, 336-337 Slide 85 Spleen, 173-175 Slide 182 Seminal Vesicle, 338-339 Slide 86 Scalp, 184-187 Slide 183 Prostate, 339-340 Slide 87 Scalp, 188 Slide 184 Penis, 340-341 Slide 88 Hair Follicle, 186 Slide 185 Epididymis, 336 Slide 88 thin skin, 189 Slide 207 Cervix, 324 Slide 90 Thin Skin, 26, 189 Slide 211 Pancreas, 221 Slide 91 Thick Skin, 27, 103-105, 190-192 Slide 214 Urinary Bladder, 287 Slide 94 Parotid Gland, 199-201 Slide 218 Mammary Gland, 327 Slide 95 Submandibular Gland, 201-203 Slide 224 Uterine Tube, 323 Slide 96 Sublingual Gland, 203-204 Slide 226 Eye, 352 Slide 109 Esophagus, 26, 96-97, 230-232 Slide 227A Eve. 348-350 Slide 110 Trachea, 298 Slide 227B Eye, 353 Slide 229 Eve, 351-352 Slide 119 Simple columnar epithelium, 23-24 Slide 120 Simple columnar epithelium, 24 Slide 230 Ear, 356-358 Slide 123 Colon, 97-98, 243-245 Slide 231 Eyelid, 354-355

Slide 125 Liver (6 mos. Fetus), 257-258